

TYENNE® (tocilizumab-aazg) Enrollment and Prescription Form

Required fields are marked with an asterisk (*).

1. Fax to KabiCare at 1-833-671-1010

2. Upload completed form file to kabicare.trialcard.com

 Fax to patient's preferred specialty pharmacy Questions? Call 1.833.KABICARE (1-833-522-4227)

	Patient informat	1011 10 00 0	ompreted by t	a patient of regu	in, additionized po		,	· ·		
	First Name*:	Last N	ame*:		D.	O.B.* /	/	Gender:	М	F
	Address*:	ddress*: City*:				9	State*:	Zip Code*:		
	Phone Number*:	one Number*: Home Cell Email*:						tion services	Υ	N
	Parent/Guardian Name	C	aregiver Nan	ne (if applicable)			needed: Language:			
	Best time to call (Monday-Friday):	Anytime	Morning	Afternoon	Evening	Check fo	or permission	to leave messages w	ith caregi	iver
	When did you start on treatment?*:	Not yet star	rted 0-3	months ago	4-6 months ago	7-12 m	onths ago	12+ month	is ago	
	Required Patient Authorization a	nd Addition	al Consents	* (all fields mus	st be completed p	orior to subr	mitting to	avoid delay in e	enrollm	ent)
	I have read and agree to the Patient Authorization to Share Personal Health Information and TYENNE® enrollment. (Section 1). I have read and agree to receive text messages and calls as explained in the Telephone Consumer Protection Act (TCPA) Consent (see section 2). I have read and agree to the Bridge to Commercial Coverage (see section 4 if applicable) and/or PAP if applicable (see section 4). I have read and agree to the Fair Credit Reporting Act (FCRA) Authorization (see section 3). Patient or authorized representative signature: Date of signature: / / If authorized representative signature, explain authority to act on behalf of patient:									
	Patient's Insurance Infor	mation - Pro	vide copies c	of the front and	back of all medic	al and presc	ription in	surance cards.		
	Beneficiary/Cardholder Name:				Prescription Ins	surance:				
2	Medical Insurance:				Rx Group #: Rx ID #:					
	Medical Insurance ID #: Group #:				Rx Bin #:		Rx P	CN #:		
	Medical modulate 15 #.				Commercial	Medica	re/ Medicaio	d VA	Unin	sured
		▼ F0	R HEALTH	CARE PROVI	DERS ONLY ▼					
	Prescriber's Name (First, Last)*: Prescriber's Address:									
٦	Office Phone:	Contact Nan	ne:		City:		State:	Zip Code:		
5	Contact Email:	C	office Fax*:		Tax ID:	;		PTAN*:		
	Provider NPI #*:	S	OC/Infusion C	Center:		Spe	cialty:	Rheumatology	0	ther
			Clini	ical Information	on					
	Diagnosis Date: / /		Clini			s:				
	Diagnosis Date: / / Prior Therapies:		Clini	Concon	on nitant Medications llergies:	:				
4			Clin i Giant cell a	Concon Drug A	nitant Medications		y ICD-10 c	ode*:		
4	Prior Therapies: Check all Rheumatoid arthritis indications Polyarticular juvenile idio		Giant cell ai Systemic ju	Concon Drug A rteritis venile idiopathic ar	nitant Medications llergies: thritis in		y ICD-10 c	ode*:		
4	Prior Therapies: Check all Rheumatoid arthritis indications Polyarticular juvenile idio in patients 2 years of age	and older	Giant cell a Systemic ju patients 2 y	Concon Drug A rteritis venile idiopathic ar rears of age and old	nitant Medications llergies: thritis in ler	Primar		ode*:		
4	Prior Therapies: Check all indications that apply: Rheumatoid arthritis Polyarticular juvenile idio in patients 2 years of age Pharmacy Pro	and older	Giant cell a Systemic ju patients 2 y	Concon Drug A rteritis venile idiopathic ar rears of age and old	nitant Medications llergies: thritis in	Primar		ode*:		
4	Prior Therapies: Check all Rheumatoid arthritis Polyarticular juvenile idio in patients 2 years of age Pharmacy Propagation of the patient's preferred specialty pharmacy:	and older escription - So	Giant cell al Systemic ju patients 2 y elect medicat	Concon Drug A rteritis venile idiopathic ar rears of age and old	nitant Medications llergies: thritis in ler	Primar		ode*:		
4	Prior Therapies: Check all indications that apply: Pharmacy Propagation of the propagati	and older escription - So	Giant cell a Systemic ju patients 2 y	Concon Drug A rteritis venile idiopathic ar years of age and old ion, fill out and s	nitant Medications llergies: thritis in der sign correspondin	Primar		ode*:		
4	Prior Therapies: Check all indications that apply: Pharmacy Properties Therapy IV Dosing Guidance: Adult Rheumatoid arthritis Polyarticular juvenile idio in patients 2 years of age Pharmacy Properties Therapy Patient's Therapy Patient's Therapy Patient's Company of the properties of the pro	and older escription - So	Giant cell al Systemic ju patients 2 y elect medicat	Concon Drug A rteritis venile idiopathic ar years of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A	nitant Medications llergies: thritis in der sign correspondin sing Guidance: rthritis (RA):	Primar ng prescripti	on below.	_		
4	Prior Therapies: Check all indications that apply: Pharmacy Properties Therapy Patient's preferred specialty pharmacy: Patient's Therapy Patient's V IV Dosing Guidance: Adult Rheumatoid Arthritis (RA): Initial 4 mg per kg every 4 weeks Increase to 8 mg per kg every 4 weeks based on compare the second of	and older escription - So weight:	Giant cell al Systemic ju patients 2 y elect medicat	Concon Drug A rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • (100 kg: 162 mg week based on	nitant Medications llergies: thritis in ler sign correspondin sing Guidance: rthritis (RA): administered subcutane clinical response	Primar og prescription eously every othe	on below.	_	ery	
4	Prior Therapies: Check all indications that apply: Patient's preferred specialty pharmacy: Patient's Therapy IV Dosing Guidance: Adult Rheumatoid arthritis Rheumatoid arthritis Polyarticular juvenile idio in patients 2 years of age Pharmacy Pr Patient's Therapy Patient's Therapy IV Dosing Guidance: Adult Rheumatoid Arthritis (RA): Initial 4 mg per kg every 4 weeks	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug A rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • (100 kg: 162 mg week based on	nitant Medications llergies: thritis in der sign correspondin sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane	Primar og prescription eously every othe	on below.	_	ery	
4	Prior Therapies: Check all indications that apply: Pharmacy Propagation of the propagati	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug A rteritis venile idiopathic ar years of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • ±100kg: 162 mg Giant Cell Arteritis (nitant Medications llergies: thritis in der sign correspondin sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane	Primar ng prescription eously every othe ously every week	on below.	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Platient's preferred specialty pharmacy Properties Therapy Patient's Therapy Patient'	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar years of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A *100 kg: 162 mg week based on *2100kg: 162 mg Giant Cell Arteritis (*162 mg given or glucocorticoids Polyarticular Juveni	thritis in der sign corresponding Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): ince every week as a subcle Idiopathic Arthritis (RI) le Idiopathic Arthritis (RI)	Primar og prescription eously every othe ously every week cutaneous injection	on below.	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Playarticular juvenile idion in patients 2 years of age Pharmacy Pr Patient's preferred specialty pharmacy: Patient's Therapy Patient's V IV Dosing Guidance: Adult Rheumatoid Arthritis (RA): Initial 4 mg per kg every 4 weeks Increase to 8 mg per kg every 4 weeks based on color Cell Arteritis (GCA): 6 mg per kg every 4 weeks in combination with a very comparation of the co	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • 2100kg: 162 mg Giant Cell Arteritis (• 162 mg given or glucocorticoids Polyarticular Juveni • 30 kg: 162 mg • 230 kg: 162 mg	thritis in der sign corresponding sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): tice every week as a subcutane guidance of the subcutane GCA: tille Idiopathic Arthritis (Fonce every three weeks once every two weeks once every two weeks once every two weeks once every two weeks	Primar og prescription cously every other ously every week cutaneous injection	on below.	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Playericular juvenile idion in patients 2 years of age Pharmacy Properties Therapy Patient's Therapy Pa	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • ≥100kg: 162 mg Giant Cell Arteritis (• 162 mg given or glucocorticoids Polyarticular Juveni • 30 kg: 162 mg Systemic Juvenile ii • ⟨30 kg: 162 mg	thritis in der sign corresponding Guidance: rthritis (RA): administered subcutane (clinical response administered subcutane GCA): cce every week as a subcutane (GCA): cce every three weeks once every three weeks diopathic Arthritis (SJIA) every two weeks	Primar og prescription cously every other ously every week cutaneous injection	on below.	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Pharmacy Propagation of the propagat	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Al rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • ≥100kg: 162 mg Giant Cell Arteritis (• 162 mg given or • 162 mg given or • 30 kg: 162 mg Systemic Juvenile Iv • 30 kg: 162 mg Systemic Juvenile Iv • 30 kg: 162 mg • ≥30 kg: 162 mg • ≥30 kg: 162 mg	thritis in der sign corresponding Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): ace every week as a subcutane decevery two weeks diopathic Arthritis (RJIA) every two weeks every week	Primar og prescription eously every other ously every week cutaneous injection PJIA):	on below.	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Playericular juvenile idion in patients 2 years of age Pharmacy Properties Therapy Patient's Therapy Pa	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • ≥ 100kg: 162 mg ion fill arteritis of • 162 mg given on • 162 mg given on • 30 kg: 162 mg > ≥ 30 kg: 162 mg Systemic Juvenile le • 30 kg: 162 mg > ≥ 30 kg: 162 mg TYENNE® s	thritis in der sign corresponding Guidance: sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): ce every week as a subcutane GCA thritis (Romane every three weeks once every three weeks once every two weeks diopathic Arthritis (SJIA every two weeks every week ubcutaneous (SC	Primar og prescription cously every other ously every week cutaneous injection PJIA): A):	on below. r week, followe	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Patient's preferred specialty pharmacy: Patient's Therapy Patient's Ther	and older escription - So weight:	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • ≥ 100kg: 162 mg ion fill arteritis of • 162 mg given on • 162 mg given on • 30 kg: 162 mg > ≥ 30 kg: 162 mg Systemic Juvenile le • 30 kg: 162 mg > ≥ 30 kg: 162 mg TYENNE® s	thritis in der sign corresponding Guidance: sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): nce every week as a subcutane devery week as a subcutane dipotation arthritis (Ropathic Arthritis (R	Primar og prescription cously every other cously every week cutaneous injection PJIA): A):) self-injecta ijector Injecta	on below. r week, followe	ed by an increase to ev		
5	Prior Therapies: Check all indications that apply: Patient's preferred specialty pharmacy: Patient's Therapy Patient	escription - Some	Giant cell ai Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar years of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A * 100 kg: 162 mg week based on * ≥100kg: 162 mg illocorficoids Polyarticular Juveni * 30 kg: 162 mg Systemic Juvenile Iv * 30 kg: 162 mg Systemic Juvenile Iv * 30 kg: 162 mg TYENNE® S Prefilled Choose frequ Once a w	thritis in der sign corresponding sing Guidance: rthritis (RA): administered subcutane (Linical response administered subcutane (GCA): acce every week as a subcutane (GCA): acce every three weeks once every three weeks diopathic Arthritis (SJIA) every two weeks every week ubcutaneous (SC I syringe Autoin ency: reek	Primar og prescription cously every other cously every week cutaneous injection PJIA): A):) self-injecta ijector Injecta	on below. r week, followers on, in combinate able t 162 mg Dispense: 1 month	ed by an increase to ev		
4	Prior Therapies: Check all indications that apply: Patient's preferred specialty pharmacy: Patient's Therapy Patient's Weeks Increase to 8 mg per kg every 4 weeks Increase to 8 mg per kg every 4 weeks based on collant Cell Arteritis (GCA): 6 mg per kg every 4 weeks in combination with a religional properticular Juvenile Idiopathic Arthritis (PJIA): 30 kg: 10 mg per kg every 4 weeks 30 kg: 8 mg per kg every 4 weeks Systemic Juvenile Idiopathic Arthritis (SJIA): 30 kg: 12 mg per kg every 2 weeks TYENNE® intravenous (IV) infusion SIG: Infuse: mg Choose frequency: Once every 2 weeks Once every 4 weeks	and older escription - So weight: clinical response tapering course of o	Giant cell a Systemic ju patients 2 y elect medicat kg	Concon Drug Ai rteritis venile idiopathic ar vears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A • 100 kg: 162 mg week based on • ≥100kg: 162 mg ion 162 mg given or glucocorticoids Polyarticular Juveni • 30 kg: 162 mg > 30 kg: 162 mg Systemic Juvenile • 430 kg: 162 mg • ≥30 kg: 162 mg TYENNE® S Prefilled Choose frequ Once a w Once eve	thritis in der sign corresponding sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): ace every week as a subcutane dipolatic Arthritis (Ropathic Arthritis (R	Primar og prescription cously every other cously every week cutaneous injection PJIA): A):) self-injecta ijector Injecta	on below. r week, follower on, in combinate able t 162 mg Dispense:	ed by an increase to ev	urse of	mes
5	Prior Therapies: Check all indications that apply: Patient's preferred specialty pharmacy: Patient's Therapy Patient's Therapy IV Dosing Guidance: Adult Rheumatoid Arthritis (RA): Initial 4 mg per kg every 4 weeks Increase to 8 mg per kg every 4 weeks based on c Giant Cell Arteritis (GCA): 6 mg per kg every 4 weeks in combination with a Polyarticular Juvenile Idiopathic Arthritis (PJIA): 30 kg: 10 mg per kg every 4 weeks >30 kg: 8 mg per kg every 4 weeks Systemic Juvenile Idiopathic Arthritis (SJIA): 30 kg: 12 mg per kg every 2 weeks >30 kg: 8 mg per kg every 2 weeks TYENNE® intravenous (IV) infusion SIG: Infuse: mg Choose frequency: Once every 2 weeks Once every 4 weeks Other:	weight: weight: dilinical response tapering course of	Giant cell al Systemic ju patients 2 y elect medicat kg lucocorticoids	Concon Drug Ai rteritis venile idiopathic ar /ears of age and old ion, fill out and s Subcutaneous Do Adult Rheumatoid A	thritis in der sign corresponding Guidance: sing Guidance: rthritis (RA): administered subcutane clinical response administered subcutane GCA): ce every week as a subcutane dipolar thritis (Florice every three weeks once every two weeks diopathic Arthritis (SJIA every two weeks every week ubcutaneous (SC I syringe Autoin ency: evek every weeks information provided the patient with a descriptions.	Primar og prescription cously every other ously every week cutaneous injection pula): A): is accurate to to cription of the least	r week, followers able t 162 mg Dispense: 1 month 2 months 3 months the best of m KabiCare pat	Refill: ny knowledge. I certifient support progra	tinify that I	am horize





TYENNE® (tocilizumab-aazg) Enrollment and Prescription Form

- 1. Fax to KabiCare at 1-833-671-1010
- 2. Upload completed form file to kabicare.trialcard.com
- 3. Fax to patient's preferred specialty pharmacy Questions? Call 1.833.KABICARE (1-833-522-4227)

Important Safety Information

RISK OF SERIOUS INFECTIONS:

Patients treated with TYENNE® (tocilizumab-aazg) are at increased risk for developing serious infections that may lead to hospitalization or death, including tuberculosis (TB), bacterial, invasive fungal, viral, or other opportunistic infections. If a serious infection develops, interrupt TYENNE until the infection is controlled.

Reported infections include:

- Active tuberculosis, which may present with pulmonary or extrapulmonary disease. Patients should be tested for latent tuberculosis before TYENNE use and during therapy. Treatment for latent infection should be initiated prior to TYENNE
- Invasive fungal infections, including candidiasis, aspergillosis, and pneumocystis.
 Patients with invasive fungal infections may present with disseminated, rather than localized, disease.
- Bacterial, viral and other infections due to opportunistic pathogens.

The risks and benefits of treatment with TYENNE should be carefully considered prior to initiating therapy in patients with chronic or recurrent infection.

Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with TYENNE, including the possible development of tuberculosis in patients who tested negative for latent tuberculosis infection prior to initiating therapy.

CONTRAINDICATION

WARNINGS AND PRECAUTIONS

Gastrointestinal Perforations

Events of gastrointestinal (GI) perforation have been reported in clinical trials, primarily as complications of diverticulitis in patients treated with tocilizumab. Use TYENNE with caution in patients who may be at increased risk for GI perforation. Promptly evaluate patients presenting with new-onset abdominal symptoms for early identification of GI perforation.

Hepatotoxicity

Serious cases of hepatic injury have been observed in patients taking intravenous or subcutaneous tocilizumab products. Some of these cases have resulted in liver transplant or death. Time to onset for cases ranged from months to years after treatment initiation. Most cases presented with marked elevations of transaminases (> 5 times ULN), and some cases presented with signs or symptoms of liver dysfunction and only mildly elevated transaminases.

Treatment with tocilizumab was associated with a higher incidence of transaminase elevations; increased frequency and magnitude of these elevations were observed when tocilizumab was used in combination with potentially hepatotoxic drugs (e.g., methotrexate).

It is not recommended to initiate TYENNE treatment in RA, GCA, PJIA, and SJIA patients with elevated transaminases ALT or AST greater than 1.5x ULN. In patients who develop elevated ALT or AST greater than 5x ULN discontinue TYENNE.

Measure liver tests promptly in patients who report symptoms that may indicate liver injury. If the patient is found to have abnormal liver tests, TYENNE treatment should be interrupted. TYENNE should only be restarted in patients with another explanation for the liver test abnormalities after normalization of the liver tests.

Laboratory Parameters

Laboratory monitoring is recommended due to potential consequences of treatment-related laboratory abnormalities in neutrophils, platelets, lipids, and liver function tests. Dosage modifications may be required.

Neutropenia: Treatment with tocilizumab products was associated with a higher incidence of neutropenia. It is not recommended to initiate TYENNE treatment in RA, GCA, PJIA, and SJIA patients with a low neutrophil count i.e., absolute neutrophil count (ANC) less than 2000 per mm³. In patients who develop an ANC less than 500 per mm³ treatment is not recommended.

Thrombocytopenia: Treatment with tocilizumab products was associated with a reduction in platelet counts. It is not recommended to initiate TYENNE in RA, GCA, PJIA, and SJIA patients with a platelet count below 100,000 per mm³. In patients who develop a platelet count less than 50,000 per mm³, treatment is not recommended.

Elevated Liver Enzymes: It is not recommended to initiate TYENNE treatment in patients with elevated transaminases ALT or AST >1.5x ULN. In patients who develop elevated ALT or AST >5x ULN, treatment is not recommended.

Lipid Abnormalities: Treatment with tocilizumab products was associated with increases in lipid parameters such as total cholesterol, triglycerides, LDL cholesterols, and/or HDL cholesterol.

Immunosuppression

The impact of treatment with tocilizumab products on the development of malignancies is not known, but malignancies were observed in clinical studies with tocilizumab. TYENNE is an immunosuppressant, and treatment with immunosuppressants may result in an increased risk of malignancies.

Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis, have been reported in association with tocilizumab products and anaphylactic events with a fatal outcome have been reported with intravenous infusion of tocilizumab products. TYENNE for intravenous use should only be infused by a healthcare professional with appropriate medical support to manage anaphylaxis. For TYENNE subcutaneous injection, advise patients to seek immediate medical attention if they experience any symptoms of a hypersensitivity reaction. If anaphylaxis or other hypersensitivity reaction occurs, stop administration of TYENNE immediately and discontinue TYENNE permanently. Do not administer TYENNE to patients with known hypersensitivity to tocilizumab products.

Demyelinating Disorders

The impact of treatment with tocilizumab products on demyelinating disorders is not known, but multiple sclerosis and chronic inflammatory demyelinating polyneuropathy were reported rarely in clinical studies. Monitor patients for signs and symptoms of demyelinating disorders. Prescribers should exercise caution in considering the use of TYENNE in patients with preexisting or recent-onset demyelinating disorders.

Active Hepatic Disease and Hepatic Impairment

Treatment with TYENNE is not recommended in patients with active hepatic disease or hepatic impairment.

Vaccinations

Avoid use of live vaccines concurrently with TYENNE. No data are available on the secondary transmission of infection from persons receiving live vaccines to patients receiving TYENNE or on the effectiveness of vaccination in patients receiving TYENNE. Patients should be brought up to date on all recommended vaccinations prior to initiation of TYENNE therapy, if possible.

ADVERSE REACTIONS

Most common adverse reactions (incidence of at least 5%): upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased ALT, injection site reactions.

DRUG INTERACTIONS

In GCA patients, no effect of concomitant corticosteroid on to cilizumab exposure was observed.

Cytochrome P450s in the liver are down-regulated by infection and inflammation stimuli including cytokines such as IL-6. Inhibition of IL-6 signaling in RA patients treated with tocilizumab products may restore CYP450 activities to higher levels than those in the absence of tocilizumab products leading to increased metabolism of drugs that are CYP450 substrates.

Exercise caution when coadministering TYENNE with CYP3A4 substrate drugs where decrease in effectiveness is undesirable, e.g., oral contraceptives, lovastatin, atorvastatin, etc.

USE IN PREGNANCY

The limited available data with tocilizumab products in pregnant women are not sufficient to determine whether there is a drug-associated risk for major birth defects and miscarriage.

You may report side effects to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Fresenius Kabi at (800) 551-7176.

Please see additional Important Safety Information in full Prescribing Information, including Boxed Warning.

INDICATIONS

TYENNE is indicated for the treatment of adult patients with moderately to severely active rheumatoid arthritis (RA) who have had an inadequate response to one or more Disease-Modifying Anti-Rheumatic Drugs (DMARDs).

TYENNE is indicated for the treatment of giant cell arteritis (GCA) in adult patients. TYENNE is indicated for the treatment of active polyarticular juvenile idiopathic arthritis (PJIA) in patients 2 years of age and older.

TYENNE is indicated for the treatment of active systemic juvenile idiopathic arthritis (SJIA) in patients 2 years of age and older.

Click to see full Prescribing Information, including Boxed Warning, for TYENNE® (tocilizumab-aazq).





TYENNE® (tocilizumab-aazg) **Enrollment and Prescription Form**

- 1. Fax to KabiCare at 1-833-671-1010
- 2. Upload completed form file to kabicare.trialcard.com
- 3. Fax to patient's preferred specialty pharmacy Questions? Call 1.833.KABICARE (1-833-522-4227)

ADDITIONAL TERMS AND CONDITIONS FOR ENROLLMENT

Section 1: Patient Authorization for the Use and Disclosure of Protected Health Information

PATIFNT AUTHORIZATION AND RELEASE TO COLLECT, USE, AND DISCLOSE HEALTH INFORMATION

By my signature above, I agree to allow my doctors, pharmacies, specialty pharmacy(ies), and health insurers (collectively "Healthcare Providers"), to use and disclose my personal health information related to this enrollment form or my use or potential use of TYENNE to Fresenius Kabi and its agents, authorized representatives, and contractors, including, without limitation, its HUB provider, as further described below.

Information to Be Disclosed: My protected health information (as such term is defined in the Health Insurance Portability and Accountability Act of 1996 ("HIPAA") and regulations thereunder), as well as other state and/or federally protected personal information, including my personal contact and other demographic information, all medical records and financial information and information relating to my treatment, the coordination of my treatment, and the delivery, packaging, and receipt of certain medication prescribed to me (collectively, my "Information").

Persons to Whom My Information May Be Disclosed: Fresenius Kabi and the KabiCare Patient Support Program, including any third parties responsible for the administration of the KabiCare Patient Support Program.

Purposes for Which the Disclosures Are to Be Made: Disclosures of my Information may be made to KabiCare Patient Support Program so that KabiCare Patient Support Program may use and disclose my Information

- purposes of: Communicating with my Healthcare Providers Oromunicating with my Heatincare Providers about my prescription and medical condition, including to facilitate the order, fulfillment and delivery of my prescription as needed;
 Establishing my eligibility for benefits from my health plan or other programs;
 Contacting my insurer on my behalf to determine if I am eligible for health insurance coverage or other funds;
 Providing appropriate product and reimbursement sunnort.

- 4) Providing appropriate product and reimbursement support;
 5) Contacting me regarding case management and/ or educational information or training offered by or through the KabiCare Patient Support Program;
 6) Contacting me regarding this Authorization or my use or potential use of my prescriptions and providing me with related communications, including through messages left for me that disclose that I take or may take certain prescription medications.
- messages left for me that disclose that I aak or may take certain prescription medications; 7) Contacting me to administer, evaluate, and improve the KabiCare Patient Support Programs, including analyzing the usage patterns and the effectiveness of services and helping to develop new products, services, and programs, and for other general business and administrative purposes; and 8) Disclosing my Information to third parties if required by law.

By signing this Authorization, I acknowledge my

- understanding that:
 I understand that I may refuse to sign this
- Indestanding triat.

 I understand that I may refuse to sign this Authorization and that my refusal to sign this Authorization will not affect my right to treatment or payment of benefits for health care. I understand that if I refuse to sign, I will not be eligible to receive support through the KabliCare Patient Support Program.

 I have the right to revoke this Authorization at any time by mailing 2250 Perimeter Park Dr., Suite 300 Morrisville, NC 27560. Revoking this Authorization will prohibit further uses and disclosures of my Information by the KabliCare Patient Support Program, except to the extent those uses and disclosures have been made in reliance on this Authorization and as permitted by applicable law.

 Certain pharmacy providers or other Healthcare
- by applicable law. Certain pharmacy providers or other Healthcare Providers may receive remuneration for the use or disclosure of my Information, as permitted by
- or disclosure of my Information, as permitted by this Authorization. Once my Information is released to KabiCare Patient Support Program based on this Authorization, my Information may not be subject to all of the protections and safequards provided by HIPAA or other federal and state privacy laws. However, I understand that KabiCare Patient Support Program has agreed to use or disclose my Information received only for the purposes described in this Authorization or as required by law. This Authorization will remain in effect for a period of two (2) years after I sign it unless a

or as required by faw.

This Authorization will remain in effect for a period of two (2) years after I sign it unless a shorter period is required by state law or is revoked by me earlier in writing.

I am entitled to receive a copy of this signed Authorization.

The patient, or the patient's authorized representative, MUST sign this form to participate in the KabiCare Patient Support Program. My signature, signifying my agreement with this Authorization, is provided on page I of this Enrollment Form, where it states, "I have read and agree to the Patient Authorization for the Use and Disclosure of Protected Health Information (section 1). If an authorized representative signs for the patient, please indicate relationship to the patient.

Section 2: Telephone Consumer Protection Act (TCPA) Consent

By checking the first box in the "Required Patient Authorization and Additional Consents" section on page 1, you understand that the personal information you provide will be shared with KabiCare, and its third-party partners, including TrialCard Incorporated. You also authorize KabiCare, and its partners, to communicate with you about

products, health conditions, copay and financial assistance. You are providing consent that you can be contacted by mail, email, phone or autodialed text messages under the guidelines of the Telephone Consumer Protection Act at the phone number you provide. You may opt out of individual communications from KabiCare or its partners individual communications from KabiCare or its partners entirely at any time by calling 1-833-KabiCare. KabiCare and companies providing services to KabiCare will not sell or rent your personally identifiable information, as described in the Privacy Policy. For clarity, phone contact includes telephonic contact for the purposes of advertising or telemarketing and may include either or both of (1) autodialed prerecorded voice calls, and (2) autodialed artificial voice calls.

You understand that you are not required to agree to receive any phone, mail, or text messages as a condition o participation in this Program, that message and data rate may apply depending on your phone carrier, and you may stop receiving text messages at any time by texting the word STOP in response to a text received from KabiCare or its partners.

Section 3: KabiCare Copay Assistance Program TERMS & CONDITIONS

To receive benefits under the Copay Assistance Program, the patient may contact the KabiCare Patient Support Program for current Program Product(s) subject to these Terms and Conditions. By participating in the Copay Assistance Program, patient acknowledges and agrees that he/she is eligible to participate and that he/she understands and agrees to comply with these Terms and Conditions.

- noditions.

 Patient must be prescribed the Program Product for an FDA-approved indication.

 Patient must have commercial (private or non-governmental) health insurance that provides coverage for the cost of the Program Product under a pharmacy or medical benefit nian
- plan.
 The Copay Assistance Program is valid for patients who have a valid prescription for a Fresenius Kabi medication and who are not reimbursed for the entire cost of the prescription by their commercial insurance plan. The Copay Assistance Program is not valid for patients enrolled in Medicaid, Medicare (including a Medicare Part D or Medicare Advantage plan, a Medigap plan, or an employer-sponsored health plan or prescription drug benefit program for Medicare-eligible retirees), Veteran Affairs health clan or programs, Department of Defense for Medicarc-eliqible retirees), Veteran Affairs levalth care programs, Department of Deferse health care programs, TRICAPE, CHAMPUS, Puerto Rico Government Health Insurance Plan, or any other state or federal medical or pharmaceutical benefit program or pharmaceutical benefit program or pharmaceutical assistance program, including any state pharmaceutical assistance programs (collectively, "Government Programs"). Patients who move from commercial insurance to Government Programs will no longer be eligible to participate in the Copay Assistance Program and agree to notify the Copay Assistance Program and sachusetts, the Copay Assistance Program of Massachusetts, the Copay Assistance Program en fire officers of the Program enter of the Assistance Program expires on the earlier of: (i) the Expiration Date set forth below; (ii) the
- (i) the Expiration Date set forth below; (ii) the date an AB-rated generic equivalent becomes available for the Program Product; or (iii) January 31, 2026, absent a change in Massachusetts state law. If the patient lives in California, the Copay Assistance Program expires on the earlier of: (i) the Expiration Date set forth below; or (ii) the date an FDA approved therapeutically equivalent for the Program Product or over the counter product with the same active ingredients becomes available.

- for the Program Product or over the counter product with the same active ingredients' becomes available. Patients must have an out-of-pocket cost for the Program Product prior to the Expiration Date of the Copay Assistance Program. The benefit available under the Copay Assistance Program is milted to the amount the patients' private health insurance company indicates that the patient is obligated to pay for up to a per syringe/annual maximum. The benefit available under the Copay Assistance Program Product subject to per syringe/annual maximum. Product subject to per syringe/annual maximums based on Program Product administration date. After reaching the maximum Copay Assistance Program benefit the patient will be responsible for all remaining out-of-pocket expenses. The patient or provider may contact the Kabicare Patient Support Program for more information. This Copay Assistance Program benefit is for commercially insured patients only. Uninsured and cash paying patients may be eligible for other types of support not part of the Copay Assistance Program pharmacy or healthcare professional agree not to seek reimbursement for all, or any part, of the benefit received by the patient through the Copay Assistance Program. Participating patients and pharmacies or health care professionals are responsible for reporting receipt of Copay Assistance Program. Participating patients and pharmacies or health care professionals are responsible for reporting receipt of Copay Assistance Program benefits as may be required by law. Patient must be a resident of the United States or the Commonwealth of Puerto Rico. Product must originate and be administered to patient in the United Originate and be administered to patient in the United Originate and be administered to patient in the United Originate and be administered to patient in the United Originate and be administered to patient in the United Originate and be administered to patient in the United

- Patient must be a resident of the United States or the Commonwealth of Puerto Rico. Product must originate and be administered to patient in the United States or the Commonwealth of Puerto Rico. All information applicable to the Copay Assistance Program requested on the KabiCare.US site must be provided, and all certifications must be provided. No other purchase is necessary. The Copay Assistance Program is not insurance. It is illegal to sell, purchase, trade, counterfeit, or

- duplicate, or offer to sell, purchase, trade counterfeit, or duplicate the Copay Assistance Program card. Void if reproduced. The Copay Assistance Program is intended to comply with all applicable laws and regulations, including, without limitation, the federal Anti-Kickback Statute, its implementing regulations, and related guidance interpreting the federal Anti-Kickback Statute.
- The Copay Assistance Program is void where prohibited by law, taxed, or restricted. The Copay Assistance Program is not transferable. No
- Assistance Program is not transferable. No substitutions are permitted. The Copay Assistance Program benefit has no cash value and cannot be combined with any other Copay Assistance Program, free trial, discount, rebate, prescription savings card, or other offer. The full value of the Copay Assistance Program benefit is intended to pass entirely to the patient. No other individual or entity is entitled to receive a discount or other amount in connection with the Copay Assistance Program.
- Copay Assistance Program.
 This offer is not conditioned on any past, present, or future purchase obligation, and the Copay Assistance Program does not obligate the use of any specific
- product or provider. To the extent applicable, this offer will be accepted
- To the extent applicable, this offer will be accepted only at participating pharmacies. KabiCare reserves the right to rescind, revoke, terminate, or amend the Copay Assistance Program at any time without notice. Data related to patient's receipt of Copay Assistance Program benefits may be collected, analyzed, and shared with KabiCare, for market research and other purposes related to assessing Copay Assistance Programs. Data shared with KabiCare will be aggregated and de-identified, meaning it will be combined with data related to other Copay Assistance Program redemptions and will not identify patient. The Terms and Conditions of the Copay Assistance Program are valid for Program Product only, and
- Program are valid for Program Product only, and Fresenius Kabi reserves the right to rescind, revoke, or amend the Program without notice.

Section 4: Bridge to Commercial Coverage - Patient Attestation

KabiCare provides a limited and temporary supply of free product through the KabiCare Commercial Bridge Program ("Commercial Bridge Program") for eligible commercially insured patients when a prior authorization request has been pending with the payer for more than 7 days and when other program eligibility criteria have been satisfied.

The patient and participating providers may not seek reimbursement for any free product provided under the Commercial Bridge Program nor does the Commercial Bridge Program include payment for product administration fees.

By signing above, the patient hereby certifies to the

I certify that I am not enrolled in any Federal health care program (such as Medicare, Medicaid, Veteran Affairs health care programs, Department of Defense health care programs, TRICARE, or CHAMPUS) if I am enrolled in the Commercial Bridge Program, I certify that all information provided herein is correct and complete, to the best of my knowledge. I acknowledge that any product provided through the Commercial Bridge Program is provided on my knowledge. I acknowledge that any product provided on through the Commercial Bridge Program is provided on a complimentary basis. I certify that I will not submit or cause to be submitted any claims for payment or reimbursement for such products to any third-party payor, including any Federal health care programs. If I and rebecome in possession of such product, I understand that such product is only for me and I will not give such product to anyone else. I agree that I will not sell, trade, distribute or otherwise transfer such product.

I understand that if I am enrolled in the Commercial Bridge Program, any future changes to prescription drug coverage, or insurance coverage may affect whether I can continue to participate in the Commercial Bridge Program. I agree to contact kablicare at 1-833-KABICARE (1-833-522-4227) and tell them about any changes to my prescription drug coverage, or insurance coverage, understand there is no purchase requirement associated with assistance through by or through the Commercial Bridge Program. I understand that completing this form does not guarantee that assistance will be provided.

Section 5: Fair Credit Reporting Act (FCRA) - Patient

The information I have provided is complete and accurate and will be used to decide if I am eligible to participate in the KabiCare Patient Assistance Program (the "PAP"). I agree that submitting my application is not a guarantee that I am entitled to participate in the PAP or that the PAF agree that submitting my applications Not a Qualantee that I am entitled to participate in the PAP or that the PAP is obligated to provide me with any assistance. I understand that the PAP can revise, change, or terminate the program at any time. I authorize my healthcare providers and my health plan or insurers to give my medical and financial information to KabiCare, which administers the PAP on behalf of Fresenius Kabi, the distributor of the medicines, and to Experian, which assesses my income and ability to pay. I authorize KabiCare and its service providers to obtain credit reports about me from one or more credit reporting agencies in order to verify my information and determine my eligibility to participate in the PAP, authorize KabiCare and Experian to review my medical and financial information and to use it only to determine if I am eligible to participate in the PAP, to operate the PAP, or as otherwise required or permitted by law. I understand and agree that KabiCare and Experian may contact me directly to verify the information I have submitted or to ask for additional information or documentation to process my application.

Section 6: Prescriber Certification and Statement of Medical Necessity

Prescriber Declaration: My signature certifies that the person named on this form is my patient, I will be supervising this patient's treatment, and the information that has been provided is complete and accurate to the best of my knowledge.

I also certify that I have made the clinical judgement that any products provided through kabiCare Patient Support Program(s), including KabiCare's Patient Assistance Program (the "PAP") are medically necessary and appropriate for the patient named on this form and will be used only by that patient. I will not use any such product or prescribe, provider, furnish, or dispense any portion thereof to any other person or patient. If I am or become in possession of such medications, I will not sell, resell, offer for sale, trade, or barter such products.

in addition, I certify that no claim for payment or reimbursement for any product furnished through KabiCare Patient Support Program(s), including the PAP, will be submitted to any third-party payer, including any Federal health care program (such as Medicare, Medicaid, Veteran Affairs health care programs, Department of Defense health care programs, TRICARE, or CHAMPUS), any other health care benefit plan, payer or patient, or returned for credit. In addition, I certify that no claim for payment or

I further certify that (a) any reimbursement investigation support or assistance provided to patients through KabiCare Patient Support Program is not made in exchange, directly or indirectly, for any past, present, or future recommendation, prescription, purchase, or use of the above therapy or any other product or service for or from anyone and (b) my decision to prescribe product was based solely on my determination of medical necessity as set forth herein. I understand that completing this form does not aurantee that assistance will be provided to does not guarantee that assistance will be provided to my patient.

Prescriber Acknowledgement: Where required by applicable law, regulation, or other applicable authority, I have obtained appropriate written authorization from the patient ("Legal Permission") permitting me to use and disclose my patients' health, demographic, and other individually identifiable information, including insurance and financial information, to Fresenius Kabi, its affiliates, its program administrator, and their respective agents, service providers and field reimbursement professionals for the purpose of prescribe the patient's resurance. service providers and field reimbursement professionals for the purpose of assessing the patients' insurance coverage and eligibility for participation in the KabiCare Patient Support Program(s), providing patient support programs, copay assistance, patient assistance, and/or reimbursement support in connection with the patient's treatment with product. I maintain records of such Legal Permission consistent with applicable law.

I appoint KabiCare on my behalf, to convey this prescription to the appropriate dispensing entity, to the extent permitted under applicable state law. Special Note: Prescribers in all states must follow applicable laws for a valid prescription. For prescribers in states with official prescription form requirements, please submit an actual prescription along with this enrollment form.

Section 7: Patient Assistance Program - Patient

Section 7: Patient Assistance Program - Patient Attestation

I certify that all information provided herein is correct and complete, to the best of my knowledge. To the extent I receive amy free product through programs offered by or through the KabiCare Patient Support Programs, including the KabiCare Patient Assistance Program ("PAP"), I acknowledge that such product is provided on a complimentary basis. I certify that I will not submit or cause to be submitted any claims for payment or reimbursement Products on any third-partip payor, including the Program of the Product of the Programs payor, in the Program of the Product of the Program of them about any changes to my income, prescription drug overage, or insurance coverage. I understand there is no purchase requirement associated with assistance through by or through any KabiCare Patient Support Program(s), including the PAE1 understand that completing this form does not guarantee that assistance will be provided.

I understand that Fresenius Kabi reserves the right at any I understand that Fresenius Rab reserves the right at any time and without notice to me to modify ana/or discontinue any or all support offered by or through KabiCare Patient Support Porgarms, including modification de

